arXiv:math/0312330v1 [math.QA] 17 Dec 2003

GRADED QUANTUM GROUPS

ALEXIS VIRELIZIER

Abstract. Starting from a Hopf algebra endowed with an action of a group π by Hopf automorphisms, we construct (by a "twisted" double method) a quasitriangular Hopf π-coalgebra. This method allows us to obtain non-trivial examples of quasitriangular Hopf π-coalgebras for any finite group π and for infinite groups π such as $\mathrm{GL}_n(\Bbbk)$. In particular, we define the graded quantum groups, which are Hopf π-coalgebras for $\pi = \mathbb{C}[[h]]^l$ and generalize the Drinfeld-Jimbo quantum enveloping algebras.

Contents

Introduction 1
1. Hopf group-coalgebras 2
2. Twisted double of Hopf algebras 5
3. The case of algebras of finite groups 11
4. Example of a quasitriangular Hopf $GL_n(k)$-coalgebra 12
5. Graded quantum groups 15
6. h-adic graded quantum groups 17
References 19

Introduction

The aim of the present paper is to construct examples of quasitriangular Hopf group-coalgebras. These algebraic structures were introduced by Turaev [Tur00]. Let π be a group. The category of representations of a quasitriangular Hopf π-coalgebra is a braided π-category. Such categories are used in [Tur00] to construct 3-dimensional homotopy quantum field theory with target an Eilenberg-Mac Lane space of type $K(\pi,1)$. Moreover, Hopf π-coalgebras are used in [Vir01, Vir02b] to construct Hennings-type and Kuperberg-type invariants of flat π-bundles over link complements and over 3-manifolds.

Let π be a group. A Hopf π-coalgebra is a family $H = \{H_\alpha\}_{\alpha\in\pi}$ of algebras (over a field $\Bbbk$) endowed with a comultiplication $\Delta = \{\Delta_{\alpha,\beta}\colon H_{\alpha\beta} \to H_\alpha \otimes H_\beta\}_{\alpha,\beta\in\pi}$, a counit $\varepsilon : H_1 \to \Bbbk$, and an antipode $S = \{S_\alpha\colon H_\alpha \to H_{\alpha^{-1}}\}_{\alpha\in\pi}$ which verify some compatibility conditions. A crossing for H is a family of algebra isomorphisms $\varphi = \{\varphi_\beta\colon H_\alpha \to H_{\beta\alpha\beta^{-1}}\}_{\alpha,\beta\in\pi}$ which preserves the comultiplication and the counit, and which yields an action of π in the sense that $\varphi_\beta\varphi_{\beta'} = \varphi_{\beta\beta'}$. A crossed Hopf π-coalgebra H is quasitriangular when it is endowed with an R-matrix $R = \{R_{\alpha,\beta} \in H_\alpha \otimes H_\beta\}_{\alpha,\beta\in\pi}$ verifying some axioms (involving the crossing φ) which generalize

---

Date: February 1, 2008.
2000 *Mathematics Subject Classification.* 81R50,17B37.

the classical ones given in [Dri87]. The case $\pi = 1$ is the standard setting of Hopf algebras.

Starting from a crossed Hopf π-coalgebra $H = \{H_\alpha\}_{\alpha \in \pi}$, Zunino [Zun02] constructed a double $Z(H) = \{Z(H)_\alpha\}_{\alpha \in \pi}$ of H which is a quasitriangular Hopf π-coalgebra in which H is embedded. One has that $Z(H)_\alpha = H_\alpha \otimes \left(\oplus_{\beta \in \pi} H^*_\beta\right)$ as a vector space. Unfortunately, each component $Z(H)_\alpha$ is infinite-dimensional (unless $H_\beta = 0$ for all but a finite number of $\beta \in \pi$).

To obtain non-trivial examples of quasitriangular Hopf π-coalgebras with finite-dimensional components, we restrict ourself to a less general situation: our initial data is not any crossed Hopf π-coalgebra but a Hopf algebra endowed with an action of π by Hopf algebra automorphisms. Remark indeed that the component H_1 of a Hopf π-coalgebra $H = \{H_\alpha\}_{\alpha \in \pi}$ is a Hopf algebra and that a crossing for H induces an action of π on H_1 by Hopf automorphisms.

In this paper, starting from a Hopf algebra A endowed with an action $\phi \colon \pi \to \mathrm{Aut}_{\mathrm{Hopf}}(A)$ of a group π by Hopf automorphisms, we construct a quasitriangular Hopf π-coalgebra $D(A, \phi) = \{D(A, \phi_\alpha)\}_{\alpha \in \pi}$. The algebra $D(A, \phi_\alpha)$ is constructed in a manner similar to the Drinfeld double (in particular $D(A, \phi_\alpha) = A \otimes A^*$ as a vector space) except that its multiplication is "twisted" by the Hopf automorphism $\phi_\alpha : A \to A$. The algebra $D(A, \mathrm{id}_A)$ is the usual Drinfeld double. In general, the algebras $D(A, \phi_\alpha)$ and $D(A, \phi_\beta)$ are not isomorphic when $\alpha \neq \beta$.

This method allows us to define non-trivial examples of quasitriangular Hopf π-coalgebras for any finite group π and for infinite groups π such as $\mathrm{GL}_n(\Bbbk)$. In particular, given a complex simple Lie algebra $\mathfrak{g}$ of rank n, we define the *graded quantum groups* $\{U_h^\alpha(\mathfrak{g})\}_{\alpha \in \mathbb{C}^{*n}}$ and $\{U_h^\alpha(\mathfrak{g})\}_{\alpha \in \mathbb{C}[[h]]^n}$ which are crossed Hopf group-coalgebras. They are obtained as quotients of $D(U_q(\mathfrak{b}_+), \phi)$ and $D(U_h(b_+), \phi')$, where $\mathfrak{b}_+$ denotes the Borel subalgebra of $\mathfrak{g}$, ϕ is an action of $\mathbb{C}^{*n}$ by Hopf automorphisms of $U_q(\mathfrak{b}_+)$, and ϕ' is an action of $\mathbb{C}[[h]]^n$ by Hopf automorphisms of $U_h(\mathfrak{b}_+)$. Furthermore, the crossed Hopf $\mathbb{C}[[h]]^n$-coalgebra $\{U_h^\alpha(\mathfrak{g})\}_{\alpha \in \mathbb{C}[[h]]^n}$ is quasitriangular.

The paper is organized as follows. In Section 1, we review the basic definitions and properties of Hopf π-coalgebras. In Section 2, we define the twisted double of a Hopf algebra A endowed with an action of a group π by Hopf automorphisms. In Section 3, we explore the case $A = \Bbbk[G]$ where G is a finite group. In Section 4, we give an example of a quasitriangular Hopf $\mathrm{GL}_n(\Bbbk)$-coalgebra. Finally, we define the graded quantum groups in Sections 5 and 6.

Throughout this paper, we let π be a group (with neutral element 1) and $\Bbbk$ be a field.

1. Hopf group-coalgebras

In this section, we review some definitions and properties concerning Hopf group-coalgebras. For a detailed treatment of the theory of Hopf group-coalgebras, we refer to [Vir02a].

1.1. **Hopf π-coalgebras.** A *Hopf π-coalgebra* (over $\Bbbk$) is a family $H = \{H_\alpha\}_{\alpha \in \pi}$ of $\Bbbk$-algebras endowed with a family $\Delta = \{\Delta_{\alpha,\beta} \colon H_{\alpha\beta} \to H_\alpha \otimes H_\beta\}_{\alpha,\beta \in \pi}$ of algebra homomorphisms (the *comultiplication*) and an algebra homomorphism $\varepsilon \colon H_1 \to \Bbbk$

(the *counit*) such that, for all $\alpha, \beta, \gamma \in \pi$,

$$(\Delta_{\alpha,\beta} \otimes \mathrm{id}_{H_\gamma})\Delta_{\alpha\beta,\gamma} = (\mathrm{id}_{H_\alpha} \otimes \Delta_{\beta,\gamma})\Delta_{\alpha,\beta\gamma}, \tag{1.1}$$

$$(\mathrm{id}_{H_\alpha} \otimes \varepsilon)\Delta_{\alpha,1} = \mathrm{id}_{H_\alpha} = (\varepsilon \otimes \mathrm{id}_{H_\alpha})\Delta_{1,\alpha}, \tag{1.2}$$

and with a family $S = \{S_\alpha \colon H_\alpha \to H_{\alpha^{-1}}\}_{\alpha\in\pi}$ of $\Bbbk$-linear maps (the *antipode*) which verifies that, for all $\alpha \in \pi$,

$$m_\alpha(S_{\alpha^{-1}} \otimes \mathrm{id}_{H_\alpha})\Delta_{\alpha^{-1},\alpha} = \varepsilon\, 1_\alpha = m_\alpha(\mathrm{id}_{H_\alpha} \otimes S_{\alpha^{-1}})\Delta_{\alpha,\alpha^{-1}}, \tag{1.3}$$

where $m_\alpha \colon H_\alpha \otimes H_\alpha \to H_\alpha$ and $1_\alpha \in H_\alpha$ denote respectively the multiplication and unit element of H_α.

When $\pi = 1$, one recovers the usual notion of a Hopf algebra. In particular $(H_1, m_1, 1_1, \Delta_{1,1}, \varepsilon, S_1)$ is a Hopf algebra.

Remark that the notion of a Hopf π-coalgebra is not self-dual and that if $H = \{H_\alpha\}_{\alpha\in\pi}$ is a Hopf π-coalgebra, then $\{\alpha \in \pi \mid H_\alpha \neq 0\}$ is a subgroup of π.

A Hopf π-coalgebra $H = \{H_\alpha\}_{\alpha\in\pi}$ is said to be of *finite type* if, for all $\alpha \in \pi$, H_α is finite-dimensional (over $\Bbbk$). Note that it does not mean that $\oplus_{\alpha\in\pi} H_\alpha$ is finite-dimensional (unless $H_\alpha = 0$ for all but a finite number of $\alpha \in \pi$).

The antipode of a Hopf π-coalgebra $H = \{H_\alpha\}_{\alpha\in\pi}$ is anti-multiplicative: each $S_\alpha \colon H_\alpha \to H_{\alpha^{-1}}$ is an anti-homomorphism of algebras, and anti-comultiplicative: $\varepsilon S_1 = \varepsilon$ and $\Delta_{\beta^{-1},\alpha^{-1}} S_{\alpha\beta} = \sigma_{H_{\alpha^{-1}},H_{\beta^{-1}}}(S_\alpha \otimes S_\beta)\Delta_{\alpha,\beta}$ for any $\alpha, \beta \in \pi$, see [Vir02a, Lemma 1.1].

The antipode $S = \{S_\alpha\}_{\alpha\in\pi}$ of $H = \{H_\alpha\}_{\alpha\in\pi}$ is said to be *bijective* if each S_α is bijective. As for Hopf algebras, the antipode of a finite type Hopf π-coalgebra is always bijective (see [Vir02a, Corollary 3.7(a)]).

We extend the Sweedler notation for the comultiplication of a Hopf π-coalgebra $H = \{H_\alpha\}_{\alpha\in\pi}$ in the following way: for any $\alpha, \beta \in \pi$ and $h \in H_{\alpha\beta}$, we write $\Delta_{\alpha,\beta}(h) = \sum_{(h)} h_{(1,\alpha)} \otimes h_{(2,\beta)} \in H_\alpha \otimes H_\beta$, or shortly, if we leave the summation implicit, $\Delta_{\alpha,\beta}(h) = h_{(1,\alpha)} \otimes h_{(2,\beta)}$. The coassociativity of Δ gives that, for any $\alpha, \beta, \gamma \in \pi$ and $h \in H_{\alpha\beta\gamma}$,

$$h_{(1,\alpha\beta)(1,\alpha)} \otimes h_{(1,\alpha\beta)(2,\beta)} \otimes h_{(2,\gamma)} = h_{(1,\alpha)} \otimes h_{(2,\beta\gamma)(1,\beta)} \otimes h_{(2,\beta\gamma)(2,\gamma)}.$$

This element of $H_\alpha \otimes H_\beta \otimes H_\gamma$ is written as $h_{(1,\alpha)} \otimes h_{(2,\beta)} \otimes h_{(3,\gamma)}$. By iterating the procedure, we define inductively $h_{(1,\alpha_1)} \otimes \cdots \otimes h_{(n,\alpha_n)}$ for any $h \in H_{\alpha_1\cdots\alpha_n}$.

1.2. **Crossed Hopf π-coalgebras.** A Hopf π-coalgebra $H = \{H_\alpha\}_{\alpha\in\pi}$ is said to be *crossed* if it is endowed with a family $\varphi = \{\varphi_\beta \colon H_\alpha \to H_{\beta\alpha\beta^{-1}}\}_{\alpha,\beta\in\pi}$ of algebra isomorphisms (the *crossing*) such that, for all $\alpha, \beta, \gamma \in \pi$,

$$(\varphi_\beta \otimes \varphi_\beta)\Delta_{\alpha,\gamma} = \Delta_{\beta\alpha\beta^{-1},\beta\gamma\beta^{-1}}\varphi_\beta, \tag{1.4}$$

$$\varepsilon\varphi_\beta = \varepsilon, \tag{1.5}$$

$$\varphi_\alpha\varphi_\beta = \varphi_{\alpha\beta}. \tag{1.6}$$

It is easy to check that $\varphi_1|_{H_\alpha} = \mathrm{id}_{H_\alpha}$ and $\varphi_\beta S_\alpha = S_{\beta\alpha\beta^{-1}}\varphi_\beta$ for all $\alpha, \beta \in \pi$.

1.3. **Quasitriangular Hopf π-coalgebras.** A crossed Hopf π-coalgebra $H = \{H_\alpha\}_{\alpha\in\pi}$ is said to be *quasitriangular* if it is endowed with a family $R = \{R_{\alpha,\beta} \in H_\alpha \otimes H_\beta\}_{\alpha,\beta\in\pi}$ of invertible elements (the *R-matrix*) such that, for all $\alpha, \beta, \gamma \in \pi$

and $x \in H_{\alpha\beta}$,

$$R_{\alpha,\beta} \cdot \Delta_{\alpha,\beta}(x) = \sigma_{\beta,\alpha}(\varphi_{\alpha^{-1}} \otimes \mathrm{id}_{H_\alpha})\Delta_{\alpha\beta\alpha^{-1},\alpha}(x) \cdot R_{\alpha,\beta}, \tag{1.7}$$

$$(\mathrm{id}_{H_\alpha} \otimes \Delta_{\beta,\gamma})(R_{\alpha,\beta\gamma}) = (R_{\alpha,\gamma})_{1\beta 3} \cdot (R_{\alpha,\beta})_{12\gamma}, \tag{1.8}$$

$$(\Delta_{\alpha,\beta} \otimes \mathrm{id}_{H_\gamma})(R_{\alpha\beta,\gamma}) = [(\mathrm{id}_{H_\alpha} \otimes \varphi_{\beta^{-1}})(R_{\alpha,\beta\gamma\beta^{-1}})]_{1\beta 3} \cdot (R_{\beta,\gamma})_{\alpha 23}, \tag{1.9}$$

$$(\varphi_\beta \otimes \varphi_\beta)(R_{\alpha,\gamma}) = R_{\beta\alpha\beta^{-1},\beta\gamma\beta^{-1}}, \tag{1.10}$$

where $\sigma_{\beta,\alpha}$ denotes the flip map $H_\beta \otimes H_\alpha \to H_\alpha \otimes H_\beta$ and, for $\Bbbk$-spaces P, Q and $r = \sum_j p_j \otimes q_j \in P \otimes Q$, we set $r_{12\gamma} = r \otimes 1_\gamma \in P \otimes Q \otimes H_\gamma$, $r_{\alpha 23} = 1_\alpha \otimes r \in H_\alpha \otimes P \otimes Q$, and $r_{1\beta 3} = \sum_j p_j \otimes 1_\beta \otimes q_j \in P \otimes H_\beta \otimes Q$.

Note that $R_{1,1}$ is a (classical) R-matrix for the Hopf algebra H_1.

When π is abelian and φ is *trivial* (that is, $\varphi_\beta|_{H_\alpha} = \mathrm{id}_{H_\alpha}$ for all $\alpha, \beta \in \pi$), one recovers the definition of a quasitriangular π-colored Hopf algebra given by Ohtsuki in [Oht93].

The R-matrix always verifies (see [Vir02a, Lemma 6.4]) that, for any $\alpha, \beta, \gamma \in \pi$,

$$(\varepsilon \otimes \mathrm{id}_{H_\alpha})(R_{1,\alpha}) = 1_\alpha = (\mathrm{id}_{H_\alpha} \otimes \varepsilon)(R_{\alpha,1}), \tag{1.11}$$

$$(S_{\alpha^{-1}}\varphi_\alpha \otimes \mathrm{id}_{H_\beta})(R_{\alpha^{-1},\beta}) = R_{\alpha,\beta}^{-1} \quad \text{and} \quad (\mathrm{id}_{H_\alpha} \otimes S_\beta)(R_{\alpha,\beta}^{-1}) = R_{\alpha,\beta^{-1}}, \tag{1.12}$$

$$(S_\alpha \otimes S_\beta)(R_{\alpha,\beta}) = (\varphi_\alpha \otimes \mathrm{id}_{H_{\beta^{-1}}})(R_{\alpha^{-1},\beta^{-1}}), \tag{1.13}$$

and provides a solution of the π-colored Yang-Baxter equation:

$$\begin{aligned}(R_{\beta,\gamma})_{\alpha 23} \cdot (R_{\alpha,\gamma})_{1\beta 3} \cdot (R_{\alpha,\beta})_{12\gamma} \\ = (R_{\alpha,\beta})_{12\gamma} \cdot [(\mathrm{id}_{H_\alpha} \otimes \varphi_{\beta^{-1}})(R_{\alpha,\beta\gamma\beta^{-1}})]_{1\beta 3} \cdot (R_{\beta,\gamma})_{\alpha 23}.\end{aligned} \tag{1.14}$$

1.4. **Ribbon Hopf π-coalgebras.** A quasitriangular Hopf π-coalgebra $H = \{H_\alpha\}_{\alpha\in\pi}$ is said to be *ribbon* if it is endowed with a family $\theta = \{\theta_\alpha \in H_\alpha\}_{\alpha\in\pi}$ of invertible elements (the *twist*) such that, for any $\alpha, \beta \in \pi$,

$$\varphi_\alpha(x) = \theta_{\alpha^{-1}} x \theta_\alpha \quad \text{for all } x \in H_\alpha, \tag{1.15}$$

$$S_\alpha(\theta_\alpha) = \theta_{\alpha^{-1}}, \tag{1.16}$$

$$\varphi_\beta(\theta_\alpha) = \theta_{\beta\alpha\beta^{-1}}, \tag{1.17}$$

$$\Delta_{\alpha,\beta}(\theta_{\alpha\beta}) = (\theta_\alpha \otimes \theta_\beta) \cdot \sigma_{\beta,\alpha}((\varphi_{\alpha^{-1}} \otimes \mathrm{id}_{H_\alpha})(R_{\alpha\beta\alpha^{-1},\alpha})) \cdot R_{\alpha,\beta}. \tag{1.18}$$

Note that θ_1 is a (classical) twist of the quasitriangular Hopf algebra H_1.

1.5. **Hopf π-coideals.** Let $H = \{H_\alpha\}_{\alpha\in\pi}$ be a Hopf π-coalgebra. A *Hopf π-coideal* of H is a family $I = \{I_\alpha\}_{\alpha\in\pi}$, where each I_α is an ideal of H_α, such that, for any $\alpha, \beta \in \pi$,

$$\Delta_{\alpha,\beta}(I_{\alpha\beta}) \subset I_\alpha \otimes H_\beta + H_\alpha \otimes I_\beta, \tag{1.19}$$

$$\varepsilon(I_1) = 0, \tag{1.20}$$

$$S_\alpha(I_\alpha) \subset I_{\alpha^{-1}}. \tag{1.21}$$

The quotient $\bar{H} = \{\bar{H}_\alpha = H_\alpha / I_\alpha\}_{\alpha\in\pi}$, endowed with the induced structure maps, is then a Hopf π-coalgebra. If H is furthermore crossed, with a crossing φ such that, for any $\alpha, \beta \in \pi$,

$$\varphi_\beta(I_\alpha) \subset I_{\alpha\beta\alpha^{-1}}, \tag{1.22}$$

then so is $\bar{H}$ (for the induced crossing).

2. Twisted double of Hopf algebras

In this section, we give a method (the *twisted double*) of defining a quasitriangular Hopf π-coalgebra from a Hopf algebra endowed with an action of a group π by Hopf automorphisms.

2.1. **Hopf pairings.**

Recall that a *Hopf pairing* between two Hopf algebras A and B (over $\Bbbk$) is a bilinear pairing $\sigma\colon A \times B \to \Bbbk$ such that, for all $a, a' \in A$ and $b, b' \in B$,

$$\sigma(a, bb') = \sigma(a_{(1)}, b)\,\sigma(a_{(2)}, b'), \tag{2.1}$$

$$\sigma(aa', b) = \sigma(a, b_{(2)})\,\sigma(a', b_{(1)}), \tag{2.2}$$

$$\sigma(a, 1) = \varepsilon(a) \quad \text{and} \quad \sigma(1, b) = \varepsilon(b). \tag{2.3}$$

Note that such a pairing always verifies that, for any $a \in A$ and $b \in B$,

$$\sigma(S(a), S(b)) = \sigma(a, b). \tag{2.4}$$

(Since both σ and $\sigma \circ (S \times S)$ are the inverse of $\sigma \circ (\mathrm{id} \times S)$ in the algebra $\mathrm{Hom}_{\Bbbk}(A \times B, \Bbbk)$ endowed with the convolution product).

Let $\sigma\colon A \times B \to \Bbbk$ be a Hopf pairing. Its annihilator ideals are $I_A = \{a \in A \,|\, \sigma(a, b) = 0 \text{ for all } b \in B\}$ and $I_B = \{b \in B \,|\, \sigma(a, b) = 0 \text{ for all } a \in A\}$. It is easy to check that I_A and I_B are Hopf ideals of A and B, respectively. Recall that σ is said to be *non-degenerate* if I_A and I_B are both reduced to 0. A degenerate Hopf pairing $\sigma\colon A \times B \to \Bbbk$ induces (by passing to the quotients) a Hopf pairing $\bar{\sigma}\colon A/I_A \times B/I_B \to \Bbbk$ which is non-degenerate.

Most of Hopf algebras we shall consider in the sequel will be defined by generators and relations. The following provides us with a method of constructing Hopf pairings, see [Dae93, KRT97].

Let $\tilde{A}$ (resp. $\tilde{B}$) be a free algebra generated by elements $a_1, \dots, a_p$ (resp. $b_1, \dots, b_q$) over $\Bbbk$. Suppose that $\tilde{A}$ and $\tilde{B}$ have Hopf algebra structures such that each $\Delta(a_i)$ for $1 \le i \le p$ (resp. $\Delta(b_j)$ for $1 \le i \le q$) is a linear combination of tensors $a_r \otimes a_s$ (resp. $b_r \otimes b_s$). Given pq scalars $\lambda_{i,j} \in \Bbbk$ with $1 \le i \le p$ and $1 \le j \le q$, there is a unique Hopf pairing $\sigma : \tilde{A} \times \tilde{B} \to \Bbbk$ such that $\sigma(a_i, b_j) = \lambda_{i,j}$.

Suppose now that A (resp. B) is the algebra obtained as the quotient of $\tilde{A}$ (resp. $\tilde{B}$) by the ideal generated by elements $r_1, \dots, r_m \in \tilde{A}$ (resp. $s_1, \dots, s_n \in \tilde{B}$). Suppose also that the Hopf algebra structure in $\tilde{A}$ (resp. $\tilde{B}$) induces a Hopf algebra structure in A (resp. B). Then a Hopf pairing $\sigma : \tilde{A} \times \tilde{B} \to \Bbbk$ induces a Hopf pairing $A \times B \to \Bbbk$ if and only if $\sigma(r_i, b_j) = 0$ for all $1 \le i \le m$ and $1 \le j \le q$, and $\sigma(a_i, s_j) = 0$ for all $1 \le i \le p$ and $1 \le j \le n$.

2.2. **The twisted double construction.**

Definition-Lemma 2.1. *Let $\sigma\colon A \times B \to \Bbbk$ be Hopf pairing between two Hopf algebras A and B. Let $\phi\colon A \to A$ be a Hopf algebra endomorphism of A. Set $D(A, B; \sigma, \phi) = A \otimes B$ as a $\Bbbk$-space. Then $D(A, B; \sigma, \phi)$ has a structure of an associative and unitary algebra given, for any $a, a' \in A$ and $b, b' \in B$, by*

$$(a \otimes b) \cdot (a' \otimes b') = \sigma(\phi(a'_{(1)}), S(b_{(1)}))\,\sigma(a'_{(3)}, b_{(3)})\; aa'_{(2)} \otimes b_{(2)}b', \tag{2.5}$$

$$1_{D(A,B;\sigma,\phi)} = 1_A \otimes 1_B. \tag{2.6}$$

Moreover, the linear embeddings $A \hookrightarrow D(A,B;\sigma,\phi)$ and $B \hookrightarrow D(A,B;\sigma,\phi)$ defined by $a \mapsto a \otimes 1_B$ and $b \mapsto 1_A \otimes b$, respectively, are algebra morphisms.

Remarks 2.2. (a) Note that $D(A,B;\sigma,\mathrm{id}_A)$ is the underlying algebra of the usual quantum double of A and B (obtained by using the Hopf pairing σ).

(b) If ϕ and ϕ' are different Hopf algebra endomorphisms of A, then the algebras $D(A,B;\sigma,\phi)$ and $D(A,B;\sigma,\phi')$ are not in general isomorphic, see Remark 4.2.

Proof. Let $a,a',a'' \in A$ and $b,b',b'' \in B$. Using the fact that σ is a Hopf pairing and ϕ is a Hopf algebra endomorphism, we have that

$$\begin{aligned}
&((a\otimes b)\cdot(a'\otimes b'))\cdot(a''\otimes b'')\\
&\quad= \sigma(\phi(a'_{(1)}),S(b_{(1)}))\,\sigma(a'_{(3)},b_{(5)})\,\sigma(\phi(a''_{(1)}),S(b_{(2)}b'_{(1)}))\\
&\qquad\qquad \sigma(a''_{(3)},b_{(4)}b'_{(3)})\;aa'_{(2)}a''_{(3)}\otimes b_{(3)}b'_{(2)}b''\\
&\quad= \sigma(\phi(a'_{(1)}),S(b_{(1)}))\,\sigma(a'_{(3)},b_{(5)})\,\sigma(\phi(a''_{(1)}),S(b'_{(1)}))\,\sigma(\phi(a''_{(2)}),S(b_{(2)}))\\
&\qquad\qquad \sigma(a''_{(4)},b_{(4)})\,\sigma(a''_{(5)},b'_{(3)})\;aa'_{(2)}a''_{(3)}\otimes b_{(3)}b'_{(2)}b'',
\end{aligned}$$

and

$$\begin{aligned}
&(a\otimes b)\cdot((a'\otimes b')\cdot(a''\otimes b''))\\
&\quad= \sigma(\phi(a''_{(1)}),S(b'_{(1)}))\,\sigma(a''_{(5)},b'_{(3)})\,\sigma(\phi(a'_{(1)}a''_{(2)}),S(b_{(1)}))\\
&\qquad\qquad \sigma(a'_{(3)}a''_{(4)},b_{(3)})\;aa'_{(2)}a''_{(3)}\otimes b_{(2)}b'_{(2)}b''\\
&\quad= \sigma(\phi(a''_{(1)}),S(b'_{(1)}))\,\sigma(a''_{(5)},b'_{(3)})\,\sigma(\phi(a'_{(1)}),S(b_{(1)}))\,\sigma(\phi(a''_{(2)}),S(b_{(2)}))\\
&\qquad\qquad \sigma(a'_{(3)},b_{(5)})\,\sigma(a''_{(4)},b_{(4)})\;aa'_{(2)}a''_{(3)}\otimes b_{(3)}b'_{(2)}b''.
\end{aligned}$$

Hence the product is associative. Finally, $1_A \otimes 1_B$ is the unit element since

$$\begin{aligned}
(a\otimes b)\cdot(1\otimes 1) &= \sigma(\phi(1),S(b_{(1)}))\,\sigma(1,b_{(3)})\;a\otimes b_{(2)}\\
&= \varepsilon(S(b_{(1)}))\,\varepsilon(b_{(3)})\;a\otimes b_{(2)} \;=\; a\otimes b,
\end{aligned}$$

and

$$\begin{aligned}
(1\otimes 1)\cdot(a\otimes b) &= \sigma(\phi(a_{(1)}),S(1))\,\sigma(a_{(3)},1)\;a_{(2)}\otimes b\\
&= \varepsilon(\phi(a_{(1)}))\,\varepsilon(a_{(3)})\;a_{(2)}\otimes b \;=\; a\otimes b.
\end{aligned}$$

Finally, for any $a,a' \in A$ and $b,b' \in B$, we have that

$$\begin{aligned}
(a\otimes 1)\cdot(a'\otimes 1) &= \sigma(\phi(a'_{(1)}),S(1))\,\sigma(a'_{(3)},1)\;aa'_{(2)}\otimes 1\\
&= \varepsilon(\phi(a'_{(1)}))\,\varepsilon(a'_{(3)})\;aa'_{(2)}\otimes 1\\
&= aa'\otimes 1
\end{aligned}$$

and

$$\begin{aligned}
(1\otimes b)\cdot(1\otimes b') &= \sigma(\phi(1),S(b_{(1)}))\,\sigma(1,b_{(3)})\;1\otimes b_{(2)}b'\\
&= \varepsilon(S(b_{(1)}))\,\varepsilon(b_{(3)})\;1\otimes b_{(2)}b'\\
&= 1\otimes bb'.
\end{aligned}$$

Therefore $A \hookrightarrow D(A,B;\sigma,\phi)$ and $B \hookrightarrow D(A,B;\sigma,\phi)$ are algebra morphisms. □

In the sequel, the group of Hopf automorphisms of a Hopf algebra A will be denoted by $\mathrm{Aut}_{\mathrm{Hopf}}(A)$.

Theorem 2.3. *Let $\sigma\colon A \times B \to \Bbbk$ be Hopf pairing between two Hopf algebras A and B, and $\phi\colon \pi \to \mathrm{Aut}_{\mathrm{Hopf}}(A)$ be group homomorphism (that is, an action of π on A by Hopf automorphisms). Then the family of algebras $D(A, B; \sigma, \phi) = \{D(A, B; \sigma, \phi_\alpha)\}_{\alpha\in\pi}$ (see Definition 2.1) has a structure of a Hopf π-coalgebra given, for any $a \in A$, $b \in B$, and $\alpha, \beta \in \pi$, by*

$$\Delta_{\alpha,\beta}(a \otimes b) = (\phi_\beta(a_{(1)}) \otimes b_{(1)}) \otimes (a_{(2)} \otimes b_{(2)}), \tag{2.7}$$

$$\varepsilon(a \otimes b) = \varepsilon_A(a)\, \varepsilon_B(b), \tag{2.8}$$

$$S_\alpha(a \otimes b) = \sigma(\varphi_\alpha(a_{(1)}), b_{(1)})\, \sigma(a_{(3)}, S(b_{(3)}))\; \varphi_\alpha S(a_{(2)}) \otimes S(b_{(2)}). \tag{2.9}$$

Proof. The coassociativity (1.1) follows directly from the coassociativity of the coproducts of A and B and the fact that $\phi_{\beta\gamma} = \phi_\beta \circ \phi_\gamma$. Axiom (1.2) is a direct consequence of $\varepsilon_A \circ \phi_\alpha = \varepsilon_A$. Since $\phi_1 = \mathrm{id}_A$ and $D(A, B; \sigma, \mathrm{id}_A)$ is underlying algebra of the usual quantum double of A and B, the counit ε is multiplicative. Let us verify that $\Delta_{\alpha,\beta}$ is multiplicative. Let $a, a' \in A$ and $b, b' \in B$. On one hand we have:

$$\begin{aligned}
&\Delta_{\alpha,\beta}((a \otimes b) \cdot (a' \otimes b')) \\
&\quad= \sigma(\phi_{\alpha\beta}(a'_{(1)}), S(b_{(1)}))\, \sigma(a'_{(3)}, b_{(3)})\; \Delta_{\alpha,\beta}(aa'_{(2)} \otimes b_{(2)}b') \\
&\quad= \sigma(\phi_{\alpha\beta}(a'_{(1)}), S(b_{(1)}))\, \sigma(a'_{(4)}, b_{(4)})\; \phi_\beta(a_{(1)}a'_{(2)}) \otimes b_{(2)}b'_{(1)} \otimes a_{(2)}a'_{(3)} \otimes b_{(3)}b'_{(2)}.
\end{aligned}$$

One the other hand,

$$\begin{aligned}
&\Delta_{\alpha,\beta}(a \otimes b) \cdot \Delta_{\alpha,\beta}(a' \otimes b') \\
&\quad= (\phi_\beta(a_{(1)}) \otimes b_{(1)} \otimes a_{(2)} \otimes b_{(2)}) \cdot (\phi_\beta(a'_{(1)}) \otimes b'_{(1)} \otimes a'_{(2)} \otimes b'_{(2)}) \\
&\quad= \sigma(\phi_\alpha\phi_\beta(a'_{(1)}), S(b_{(1)}))\, \sigma(\phi_\beta(a'_{(3)}), b_{(3)})\, \sigma(\phi_\beta(a'_{(4)}), S(b_{(4)}))\, \sigma(a'_{(6)}, b_{(6)}) \\
&\qquad\qquad \phi_\beta(a_{(1)})\phi_\beta(a'_{(2)}) \otimes b_{(2)}b'_{(1)} \otimes a_{(2)}a'_{(5)} \otimes b_{(5)}b'_{(2)} \\
&\quad= \sigma(\phi_{\alpha\beta}(a'_{(1)}), S(b_{(1)}))\, \sigma(\phi_\beta(a'_{(3)}), b_{(3)}S(b_{(4)}))\, \sigma(a'_{(5)}, b_{(6)}) \\
&\qquad\qquad \phi_\beta(a_{(1)}a'_{(2)}) \otimes b_{(2)}b'_{(1)} \otimes a_{(2)}a'_{(4)} \otimes b_{(5)}b'_{(2)} \\
&\quad= \sigma(\phi_{\alpha\beta}(a'_{(1)}), S(b_{(1)}))\, \sigma(a'_{(4)}, b_{(4)})\phi_\beta(a_{(1)}a'_{(2)}) \otimes b_{(2)}b'_{(1)} \otimes a_{(2)}a'_{(3)} \otimes b_{(3)}b'_{(2)}.
\end{aligned}$$

Let us verify the first equality of (1.3). Let $a \in A$, $b \in B$, and $\alpha \in \pi$. Denote the multiplication in $D(A, B; \sigma, \phi_\alpha)$ by m_α. We have

$$\begin{aligned}
&m_\alpha(S_{\alpha^{-1}} \otimes \mathrm{id}_{D(A,B;\sigma,\phi_\alpha)})\Delta_{\alpha^{-1},\alpha}(a \otimes b) \\
&\quad= \sigma(a_{(1)}, b_{(1)})\, \sigma(\phi_\alpha(a_{(3)}), S(b_{(5)}))\, \sigma(\phi_\alpha(a_{(4)}), S^2(b_{(4)}))\, \sigma(a_{(6)}, S(b_{(2)})) \\
&\qquad\qquad S(a_{(2)})a_{(5)} \otimes S(b_{(3)})b_{(6)} \\
&\quad= \sigma(a_{(1)}, b_{(1)})\, \sigma(\phi_\alpha(a_{(3)}), S(b_{(5)})S^2(b_{(4)}))\, \sigma(a_{(5)}, S(b_{(2)})) \\
&\qquad\qquad S(a_{(2)})a_{(4)} \otimes S(b_{(3)})b_{(6)} \\
&\quad= \sigma(a_{(1)}, b_{(1)})\, \sigma(a_{(4)}, S(b_{(2)}))\; S(a_{(2)})a_{(3)} \otimes S(b_{(3)})b_{(4)} \\
&\quad= \sigma(a_{(1)}, b_{(1)})\, \sigma(a_{(2)}, S(b_{(2)}))\; 1 \otimes 1 \\
&\quad= \sigma(a, b_{(1)}S(b_{(2)}))\; 1 \otimes 1 \;=\; \varepsilon(a)\, \varepsilon(b)\; 1 \otimes 1
\end{aligned}$$

The second equality of (1.3) can be verified similarly. □

Let $\sigma\colon A \times B \to \Bbbk$ be a Hopf pairing between two Hopf algebras A and B, and $\phi\colon \pi \to \mathrm{Aut}_{\mathrm{Hopf}}(A)$ be an action of π on A by Hopf automorphisms. An action

$\psi\colon \pi \to \mathrm{Aut}_{\mathrm{Hopf}}(B)$ of π on B by Hopf automorphisms is said to be (σ, ϕ)-*compatible* if, for all $a \in A$, $b \in B$ and $\beta \in \pi$,

$$\sigma(\phi_\beta(a), \psi_\beta(b)) = \sigma(a, b). \tag{2.10}$$

Lemma 2.4. *Let $\sigma\colon A \times B \to \Bbbk$ be a Hopf pairing between two Hopf algebras A and B, and $\phi\colon \pi \to \mathrm{Aut}_{\mathrm{Hopf}}(A)$, $\psi\colon \pi \to \mathrm{Aut}_{\mathrm{Hopf}}(B)$ be two actions of π by Hopf automorphisms on A and B, respectively. Suppose that ψ is (σ, ϕ)-compatible. Then the Hopf π-coalgebra $D(A, B; \sigma, \phi) = \{D(A, B; \sigma, \phi_\alpha)\}_{\alpha \in \pi}$ (see Theorem 2.3) admits a crossing φ given, for any $a \in A$, $b \in B$ and $\beta \in \pi$, by*

$$\varphi_\beta(a \otimes b) = \phi_\beta(a) \otimes \psi_\beta(b). \tag{2.11}$$

Proof. Let $\alpha, \beta \in \pi$. We have that $\varphi_\beta(1_A \otimes 1_B) = \phi_\beta(1_A) \otimes \psi_\beta(1_B) = 1_A \otimes 1_B$ and, for any $a, a' \in A$ and $b, b' \in B$,

$$\begin{aligned}
&\varphi_\beta(a \otimes b) \cdot \varphi_\beta(a' \otimes b') \\
&\quad = \sigma(\phi_{\beta\alpha\beta^{-1}}(\phi_\beta(a')_{(1)}), S(\psi_\beta(b)_{(1)}))\, \sigma(\phi_\beta(a')_{(3)}, \psi_\beta(b)_{(3)})\ \phi_\beta(a)\phi_\beta(a')_{(2)} \otimes \psi_\beta(b)_{(2)}\psi_\beta(b') \\
&\quad = \sigma(\phi_\beta\phi_\alpha(a'_{(1)})), \psi_\beta S(b_{(1)}))\, \sigma(\phi_\beta(a'_{(3)}), \psi_\beta(b_{(3)}))\ \phi_\beta(a)\phi_\beta(a'_{(2)}) \otimes \psi_\beta(b_{(2)})\psi_\beta(b') \\
&\quad = \sigma(\phi_\alpha(a'_{(1)})), S(b_{(1)}))\, \sigma(a'_{(3)}, b_{(3)})\ \phi_\beta(aa'_{(2)}) \otimes \psi_\beta(b_{(2)}b') \\
&\quad = \varphi_\beta((a \otimes b) \cdot (a' \otimes b')).
\end{aligned}$$

Moreover ϕ_β and ψ_β are bijective and so is φ_β. Therefore $\varphi_\beta : D(A, B; \sigma, \phi_\alpha) \to D(A, B; \sigma, \phi_{\beta\alpha\beta^{-1}})$ is an algebra isomorphism.

Let $a \in A$, $b \in B$ and $\alpha, \beta, \gamma \in \pi$. We have that

$$\begin{aligned}
\Delta_{\beta\alpha\beta^{-1},\beta\gamma\beta^{-1}}(\varphi_\beta(a \otimes b)) &= \phi_{\beta\gamma\beta^{-1}}(\phi_\beta(a)_{(1)}) \otimes \psi_\beta(b)_{(1)} \otimes \phi_\beta(a)_{(2)} \otimes \psi_\beta(b)_{(2)} \\
&= \phi_{\beta\gamma\beta^{-1}}\phi_\beta(a_{(1)}) \otimes \psi_\beta(b_{(1)}) \otimes \phi_\beta(a_{(2)}) \otimes \psi_\beta(b_{(2)}) \\
&= \phi_\beta\phi_\gamma(a_{(1)}) \otimes \psi_\beta(b_{(1)}) \otimes \phi_\beta(a_{(2)}) \otimes \psi_\beta(b_{(2)}) \\
&= (\varphi_\beta \otimes \varphi_\beta)\Delta_{\alpha,\gamma}(a \otimes b),
\end{aligned}$$

$$\varepsilon\varphi_\beta(a \otimes b) = \varepsilon(\phi_\beta(a))\, \varepsilon(\psi_\beta(b)) = \varepsilon(a)\, \varepsilon(b) = \varepsilon(a \otimes b),$$

and

$$\varphi_\alpha\varphi_\beta(a \otimes b) = \phi_\alpha\phi_\beta(a) \otimes \psi_\alpha\psi_\beta(b) = \phi_{\alpha\beta}(a) \otimes \psi_{\alpha\beta}(b) = \varphi_{\alpha\beta}(a \otimes b).$$

Therefore φ satisfies Axioms (1.4), (1.5) and (1.6). □

Corrolary 2.5. *Let $\sigma\colon A \times B \to \Bbbk$ be a non-degenerate Hopf pairing and $\phi\colon \pi \to \mathrm{Aut}_{\mathrm{Hopf}}(A)$ be an action of π on A by Hopf automorphisms. Then there exists a unique action $\phi^*\colon \pi \to \mathrm{Aut}_{\mathrm{Hopf}}(B)$ which is (σ, ϕ)-compatible. It is characterized, for any $a \in A$, $b \in B$ and $\beta \in \pi$, by*

$$\sigma(a, \phi^*_\beta(b)) = \sigma(\phi_{\beta^{-1}}(a), b). \tag{2.12}$$

Consequently the Hopf π-coalgebra $D(A, B; \sigma, \phi) = \{D(A, B; \sigma, \phi_\alpha)\}_{\alpha \in \pi}$ (see Theorem 2.3) is crossed with crossing defined by $\varphi_\beta = \phi_\beta \otimes \phi^*_\beta$ for any $\beta \in \pi$.

Proof. Let $\beta \in \pi$. Since σ is non-degenerate, (2.12) does define a linear map $\phi^*_\beta\colon B \to B$. Since σ is a Hopf pairing and $\phi_{\beta^{-1}}$ is a Hopf algebra isomorphism of A, the map ϕ^*_β is a Hopf algebra isomorphism of B. Moreover ϕ^* is an action since $\phi^*_1 = \mathrm{id}_B$ (because $\phi_1 = \mathrm{id}_A$) and, for any $a \in A$, $b \in B$ and $\alpha, \beta \in \pi$,

$$\begin{aligned}
\sigma(a, \phi^*_{\alpha\beta}(b)) = \sigma(\phi_{\beta^{-1}\alpha^{-1}}(a), b) &= \sigma(\phi_{\beta^{-1}}\phi_{\alpha^{-1}}(a), b) \\
&= \sigma(\phi_{\alpha^{-1}}(a), \phi^*_\beta(b)) = \sigma(a, \phi^*_\alpha\phi^*_\beta(b)).
\end{aligned}$$

Finally (2.12) says exactly that ϕ^* is (σ, ϕ)-compatible. □

Theorem 2.6. *Let $\sigma\colon A \times B \to \Bbbk$ be a Hopf pairing between two Hopf algebras A and B, and $\phi\colon \pi \to \mathrm{Aut}_{\mathrm{Hopf}}(A)$ be an action of π on A by Hopf automorphisms. Suppose that σ is non-degenerate and that A (and so B) is finite dimensional. Then the crossed Hopf π-coalgebra $D(A, B; \sigma, \phi) = \{D(A, B; \sigma, \phi_\alpha)\}_{\alpha\in\pi}$ (see Corollary 2.5) is quasitriangular with R-matrix given, for all $\alpha, \beta \in \pi$, by*

$$R_{\alpha,\beta} = \sum_i (e_i \otimes 1_B) \otimes (1_A \otimes f_i), \tag{2.13}$$

where $(e_i)_i$ and $(f_i)_i$ are basis of A and B, respectively, such that $\sigma(e_i, f_j) = \delta_{i,j}$.

Remarks 2.7. (a) The element $\sum_i (e_i \otimes 1_B) \otimes (1_A \otimes f_i) \in A \otimes B \otimes A \otimes B$ is canonical, i.e., independent of the choices of the basis $(e_i)_i$ of A and $(f_i)_i$ of B such that $\sigma(e_i, f_j) = \delta_{i,j}$.

(b) The hypothesis A is finite dimensional is to ensure that the sum $\sum_i (e_i \otimes 1_B) \otimes (1_A \otimes f_i)$ lies in $A \otimes B \otimes A \otimes B$. More generally, assume that A and B are graded Hopf algebras with finite dimensional homogeneous components and that σ is compatible with the gradings. Then the quotient Hopf algebras A/I_A and B/I_B are also graded and can be identified via σ with the duals of each other. Suppose also that the action ϕ respects the grading so does the quotient $\bar\phi : \pi \to \mathrm{Aut}_{\mathrm{Hopf}}(A/I_A)$. In this case, there exists a unique action $\pi \to \mathrm{Aut}_{\mathrm{Hopf}}(B/I_B)$ which is $(\bar\sigma, \bar\phi)$-compatible, where $\bar\sigma\colon A/I_A \times B/I_B \to \Bbbk$ is the induced Hopf pairing. Then the Hopf π-coalgebra $D(A/I_A, B/I_B; \bar\sigma, \bar\phi)$ is quasitriangular by the same construction as in Theorem 2.6.

Proof. Fix basis (e_i) of A and (f_i) of B such that $\sigma(e_i, f_j) = \delta_{i,j}$ (such basis always exist since σ is non-degenerate). Note that $x = \sum_i \sigma(x, f_i)\, x$ and $y = \sum_i \sigma(e_i, y)\, y$ for any $x \in A$ and $y \in B$.

Recall that, since $\sum_i e_i \otimes 1_B \otimes 1_A \otimes f_i$ is the R-matrix of the usual quantum double $D(A, B, \sigma, \mathrm{id}_A)$, we have

$$\sum_{i,j} S(e_i)e_j \otimes f_i f_j = 1_A \otimes 1_B, \tag{2.14}$$

$$\sum_i e_i \otimes f_{i(1)} \otimes f_{i(2)} = \sum_{i,j} e_i e_j \otimes f_j \otimes f_i, \tag{2.15}$$

$$\sum_i e_{i(1)} \otimes e_{i(2)} \otimes f_i = \sum_{i,j} e_i \otimes e_j \otimes f_i f_j. \tag{2.16}$$

Let $\alpha, \beta \in \pi$. From (2.14) and since A (resp. B) can be viewed as a subalgebra of $D(A, B; \sigma, \phi_\alpha)$ (resp. $D(A, B; \sigma, \phi_\beta)$) via $a \mapsto a \otimes 1_B$ (resp. $b \mapsto 1_A \otimes b$), we get that $R_{\alpha,\beta}$ is invertible in $D(A, B; \sigma, \phi_\alpha) \otimes D(A, B; \sigma, \phi_\beta)$ with inverse

$$R^{-1}_{\alpha,\beta} = \sum_i S(e_i) \otimes 1_B \otimes 1_A \otimes f_i.$$

Let $a \in A$, $b \in B$ and $\alpha, \beta \in \pi$. For all $x \in A$, we have that

$$\begin{aligned}
&(\mathrm{id}_{A\otimes B\otimes A} \otimes \sigma(x,\cdot))(R_{\alpha,\beta} \cdot \Delta_{\alpha,\beta}(a \otimes b))\\
&\quad = \sum\nolimits_i \sigma(\phi_\beta(a_{(2)}), S(f_{i(1)}))\, \sigma(a_{(4)}, f_{i(3)})\, \sigma(x, f_{i(2)} b_{(2)})\; e_i \phi_\beta(a_{(1)}) \otimes b_{(1)} \otimes a_{(3)}\\
&\quad = \sum\nolimits_i \sigma(\phi_\beta S^{-1}(a_{(2)}), f_{i(1)})\, \sigma(a_{(4)}, f_{i(3)})\, \sigma(x_{(1)}, f_{i(2)}) \sigma(x_{(2)}, b_{(2)})\; e_i \phi_\beta(a_{(1)}) \otimes b_{(1)} \otimes a_{(3)}\\
&\quad = \sum\nolimits_i \sigma(a_{(4)} x_{(1)} \phi_\beta S^{-1}(a_{(2)}), f_i)\, \sigma(x_{(2)}, b_{(2)})\; e_i \phi_\beta(a)_{(1)} \otimes b_{(1)} \otimes a_{(3)}\\
&\quad = \sigma(x_{(2)}, b_{(2)})\; a_{(4)} x_{(1)} \phi_\beta(S^{-1}(a_{(2)}) a_{(1)}) \otimes b_{(1)} \otimes a_{(3)}\\
&\quad = \sigma(x_{(2)}, b_{(2)})\; a_{(2)} x_{(1)} \otimes b_{(1)} \otimes a_{(1)},
\end{aligned}$$

and, since $x_{(1)} \otimes x_{(2)} \otimes x_{(2)} \otimes x_{(2)} = \sum_i \sigma(x_{(2)}, f_i)\; x_{(1)} \otimes e_{i(1)} \otimes e_{i(2)} \otimes e_{i(3)}$,

$$\begin{aligned}
&(\mathrm{id}_{A\otimes B\otimes A} \otimes \sigma(x,\cdot))(\sigma_{\beta,\alpha}(\varphi_{\alpha^{-1}} \otimes \mathrm{id}_{H_\alpha}) \Delta_{\alpha\beta\alpha^{-1},\alpha}(a \otimes b) \cdot R_{\alpha,\beta})\\
&\quad = \sum\nolimits_i \sigma(\phi_\alpha(e_{i(1)}), S(b_{(2)}))\, \sigma(e_{i(3)}, b_{(4)})\, \sigma(x, \phi^*_{\alpha^{-1}}(b_{(1)}) f_i)\; a_{(2)} e_{i(2)} \otimes b_{(3)} \otimes a_{(1)}\\
&\quad = \sum\nolimits_i \sigma(\phi_\alpha(e_{i(1)}), S(b_{(2)}))\, \sigma(e_{i(3)}, b_{(4)})\, \sigma(\phi_\alpha(x_{(1)}), b_{(1)})\, \sigma(x_{(2)}, f_i)\; a_{(2)} e_{i(2)} \otimes b_{(3)} \otimes a_{(1)}\\
&\quad = \sigma(\phi_\alpha(x_{(2)}), S(b_{(2)}))\, \sigma(x_{(4)}, b_{(4)})\, \sigma(\phi_\alpha(x_{(1)}), b_{(1)})\; a_{(2)} x_{(3)} \otimes b_{(3)} \otimes a_{(1)}\\
&\quad = \sigma(\phi_\alpha(x_{(1)}), b_{(1)} S(b_{(2)}))\, \sigma(x_{(3)}, b_{(4)})\; a_{(2)} x_{(2)} \otimes b_{(3)} \otimes a_{(1)}\\
&\quad = \sigma(x_{(2)}, b_{(2)})\; a_{(2)} x_{(1)} \otimes b_{(1)} \otimes a_{(1)}.
\end{aligned}$$

Hence, since σ is non-degenerate, Axiom (1.7) is satisfied.

Let us verify Axioms (1.10). Let $\alpha, \beta, \gamma \in \pi$. Since ϕ^* is (σ, ϕ)-compatible (by definition), the basis $(\phi_\beta(e_i))_i$ of A and $(\phi^*_\beta(f_i))_i$ of B satisfy $\sigma(\phi_\beta(e_i), \phi^*_\beta(e_j)) = \sigma(e_i, f_j) = \delta_{i,j}$. Therefore we have that

$$(\varphi_\beta \otimes \varphi_\beta)(R_{\alpha,\gamma}) = \sum_i \phi_\beta(e_i) \otimes 1_B \otimes 1_A \otimes \phi^*_\beta(f_j) = R_{\beta\alpha\beta^{-1},\beta\gamma\beta^{-1}}.$$

Finally, let us check Axioms (1.8) and (1.9). Let $\alpha, \beta, \gamma \in \pi$. Using (2.15), we have

$$\begin{aligned}
(\mathrm{id}_{D(A,B;\sigma,\phi_\alpha)} \otimes \Delta_{\beta,\gamma})(R_{\alpha,\beta\gamma}) &= \sum\nolimits_i e_i \otimes 1_B \otimes 1_A \otimes f_{i(1)} \otimes 1_A \otimes f_{i(2)}\\
&= \sum\nolimits_{i,j} e_i e_j \otimes 1_B \otimes 1_A \otimes f_j \otimes 1_A \otimes f_i\\
&= (R_{\alpha,\gamma})_{1\beta 3} \cdot (R_{\alpha,\beta})_{12\gamma}.
\end{aligned}$$

Using (2.16) and (1.10), we have

$$\begin{aligned}
&(\Delta_{\alpha,\beta} \otimes \mathrm{id}_{D(A,B;\sigma,\phi_\gamma)})(R_{\alpha\beta,\gamma})\\
&\quad = \sum\nolimits_i \phi_\beta(e_{i(1)}) \otimes 1_B \otimes e_{i(2)} \otimes 1_B \otimes 1_A \otimes f_i\\
&\quad = \sum\nolimits_{i,j} \phi_\beta(e_i) \otimes 1_B \otimes e_j \otimes 1_B \otimes 1_A \otimes f_i f_j\\
&\quad = [(\varphi_\beta \otimes \mathrm{id}_{D(A,B;\sigma,\phi_\gamma)})(R_{\beta^{-1}\alpha\beta,\gamma})]_{1\beta 3} \cdot (R_{\beta,\gamma})_{\alpha 23}\\
&\quad = [(\mathrm{id}_{D(A,B;\sigma,\phi_\alpha)} \otimes \varphi_{\beta^{-1}})(R_{\alpha,\beta\gamma\beta^{-1}})]_{1\beta 3} \cdot (R_{\beta,\gamma})_{\alpha 23}.
\end{aligned}$$

This completes the proof of the quasitriangularity of $D(A, B; \sigma, \phi)$. □

The following corollary is a direct consequence of Corollary 2.5 and Theorem 2.6.

Corrolary 2.8. *Let A be a finite-dimensional Hopf algebra and $\phi\colon \pi \to \mathrm{Aut}_{\mathrm{Hopf}}(A)$ be an action of π on A by Hopf algebras automorphisms. Recall that the duality bracket $\langle,\rangle_{A\otimes A^*}$ is a non-degenerate Hopf pairing between A and $A^{*\mathrm{cop}}$. Then $D(A, A^{*\mathrm{cop}}; \langle,\rangle_{A\otimes A^*}, \phi)$ is a quasitriangular Hopf π-coalgebra.*

Remark 2.9. Recall that the group of Hopf automorphisms of a finite-dimensional semisimple Hopf algebra A over a field of characteristic 0 is finite (see [Rad90]). To obtain non-trivial examples of (quasitriangular) Hopf π-coalgebras for an infinite group π by using the twisted double method, one has to consider non-semisimple Hopf algebras (at least in characteristic 0).

2.3. **The h-adic case.** In this subsection, we develop the h-adic variant of Hopf group-coalgebras. A technical argument for the need of h-adic Hopf group-coalgebras is that they are necessary for a mathematically rigorous treatment of R-matrices for quantize enveloping algebras endowed with a group action.

Recall that if V is a vector space over $\mathbb{C}[[h]]$, the topology on V for which the sets $\{h^nV + v \mid n \in \mathbb{N}\}$ are a neighborhood base of $v \in V$ is called the *h-adic topology.* If V and W are vector spaces over $\mathbb{C}[[h]]$, we shall denote by $V\hat{\otimes}W$ the completion of the tensor product space $V \otimes_{\mathbb{C}[[h]]} W$ in the h-adic topology. Let V be a complex vector space. Then the set $V[[h]]$ of all formal power series $f = \sum_{n=0}^{\infty} v_n h^n$ with coefficients $v_n \in V$ is a vector space over $\mathbb{C}[[h]]$ which is complete in the h-adic topology. Furthermore, $V[[h]]\hat{\otimes}W[[h]] = (V \otimes W)[[h]]$ for any complex vector spaces V and W.

An *h-adic algebra* is a vector space A over $\mathbb{C}[[h]]$ which is complete in the h-adic topology and endowed with a $\mathbb{C}[[h]]$-linear map $m\colon A\hat{\otimes}A \to A$ and an element $1 \in A$ satisfying $m(\mathrm{id}_A\hat{\otimes}m) = m(m\hat{\otimes}\mathrm{id}_A)$ and $m(a\hat{\otimes}1) = a = m(1\hat{\otimes}a)$ for all $a \in A$.

By an *h-adic Hopf π-coalgebra*, we shall mean a family $H = \{H_\alpha\}_{\alpha\in\pi}$ of h-adic algebras which is endowed with h-adic algebra homomorphisms $\Delta_{\alpha,\beta}\colon H_{\alpha\beta} \to H_\alpha\hat{\otimes}H_\beta$ $(\alpha, \beta \in \pi)$ and $\varepsilon\colon A \to \mathbb{C}[[h]]$ satisfying (1.1) and (1.2), and with $C[[h]]$-linear maps $S_\alpha\colon H_\alpha \to H_{\alpha^{-1}}$ $(\alpha \in \pi)$ satisfying (1.3). In the previous axioms, one has to replace the algebraic tensor products $\otimes$ by the h-adic completions $\hat{\otimes}$.

The notions of crossed and quasitriangular h-adic Hopf π-coalgebras can be defined similarly as in Sections 1.2 and 1.3.

The definitions of Section 2 and Theorem 2.3 carry over almost verbatim to h-adic Hopf algebras. The only modifications are that $\sigma\colon A\hat{\otimes}B \to \mathbb{C}[[h]]$ is $\mathbb{C}[[h]]$-linear and that the algebra $D(A, B; \sigma, \phi)$, where ϕ is an h-adic Hopf endomorphism of A, is built over the completion $A\hat{\otimes}B$ of $A \otimes B$ in the h-adic topology. The reasoning of the proof of Theorem 2.6 give the following h-adic version.

Theorem 2.10. *Let $\sigma\colon A\hat{\otimes}B \to \mathbb{C}[[h]]$ be an h-adic Hopf pairing between two h-adic Hopf algebras A and B, and $\phi\colon \pi \to \mathrm{Aut}_{\mathrm{Hopf}}(A)$ be an action of π on A by h-adic Hopf automorphisms. Suppose that σ is non-degenerate. Let $(e_i)_i$ and $(f_i)_i$ be dual basis of the vector spaces A and B, respectively, with respect to the form σ. If $R_{\alpha,\beta} = \sum_i (e_i \otimes 1_B) \otimes (1_A \otimes f_i)$ belongs to the h-adic completion $D(A, B; \sigma, \phi_\alpha)\hat{\otimes}D(A, B; \sigma, \phi_\beta)$, then $R = \{R_{\alpha,\beta}\}_{\alpha,\beta\in\pi}$ is a R-matrix of the crossed h-adic Hopf π-coalgebra $D(A, B; \sigma, \phi) = \{D(A, B; \sigma, \phi_\alpha)\}_{\alpha\in\pi}$.*

3. The case of algebras of finite groups

Let G be a finite group. In this section, we describe the Hopf G-coalgebras obtained by the twisted double method from the Hopf algebra $\Bbbk[G]$.

Recall that the Hopf algebra structure of the (finite-dimensional) $\Bbbk$-algebra $\Bbbk[G]$ of G is given by $\Delta(g) = g \otimes g$, $\varepsilon(g) = 1$ and $S(g) = g^{-1}$ for all $g \in G$. The dual of $\Bbbk[G]$ is the Hopf algebra $F(G) = \Bbbk^G$ of functions $G \to \Bbbk$. It has a basis $(e_g \colon G \to \Bbbk)_{g \in G}$ defined by $e_g(h) = \delta_{g,h}$ where $\delta_{g,g} = 1$ and $\delta_{g,h} = 0$ if $g \neq h$. The structure maps of $F(G)$ are given by $e_g e_h = \delta_{g,h}\, e_g$, $1_{F(G)} = \sum_{g \in \pi} e_g$, $\Delta(e_g) = \sum_{xy=g} e_x \otimes e_y$, $\varepsilon(e_g) = \delta_{g,1}$ and $S(e_g) = e_{g^{-1}}$ for any $g, h \in G$.

Set $\phi \colon \Bbbk[G] \to \mathrm{Aut}_{\mathrm{Hopf}}(\Bbbk[G])$ defined by $\phi_g(h) = ghg^{-1}$. It is a well defined group homomorphism (since any $g \in G$ is grouplike in $\Bbbk[G]$). By Corollary 2.8, this data leads to a quasitriangular Hopf G-coalgebra $D(\Bbbk[G], F(G); \langle , \rangle_{\Bbbk[G] \times F(G)}, \phi)$ which will be denoted by $D_G(G) = \{D_\alpha(G)\}_{\alpha \in G}$.

Let us describe $D_G(G)$ more precisely. For any $\alpha \in G$, the algebra structure of $D_\alpha(G)$, which is equal to $\Bbbk[G] \otimes F(G)$ as a $\Bbbk$-space, is given by

$$(g \otimes e_h) \cdot (g' \otimes e_{h'}) = \delta_{\alpha g' \alpha^{-1}, h^{-1} g' h'}\, gg' \otimes e_{h'} \quad \text{for all } g, g', h, h' \in G,$$

$$1_{D_\alpha(G)} = \sum_{g \in G} 1 \otimes e_g.$$

The structure maps of $D_G(G)$ are given, for any $\alpha, \beta \in G$ and $g, h \in G$, by

$$\Delta_{\alpha,\beta}(g \otimes e_h) = \sum_{xy=h} \beta g \beta^{-1} \otimes e_y \otimes g \otimes e_x,$$

$$\varepsilon(g \otimes e_h) = \delta_{h,1},$$

$$S_\alpha(g \otimes e_h) = \alpha g^{-1} \alpha^{-1} \otimes e_{\alpha g \alpha^{-1} h^{-1} g^{-1}},$$

$$\varphi_\alpha(g \otimes e_h) = \alpha g \alpha^{-1} \otimes e_{\alpha h \alpha^{-1}}.$$

The crossed Hopf G-coalgebra $D_G(G)$ is quasitriangular and furthermore ribbon with R-matrix and twist given, for any $\alpha, \beta \in G$, by

$$R_{\alpha,\beta} = \sum_{g,h \in G} g \otimes e_h \otimes 1 \otimes e_g \quad \text{and} \quad \theta_\alpha = \sum_{g \in G} \alpha^{-1} g \alpha \otimes e_g.$$

Note that $\theta_\alpha^n = \sum_{g \in G} \alpha^{-n} (g\alpha)^n \otimes e_g$ for any $n \in \mathbb{Z}$.

4. EXAMPLE OF A QUASITRIANGULAR HOPF $GL_n(k)$-COALGEBRA

In this section, $\Bbbk$ is a field whose characteristic is not 2. Fix a positive integer n. We use a (finite dimensional) Hopf algebra whose group of automorphisms is known to be the group $\mathrm{GL}_n(\Bbbk)$ of invertible $n \times n$-matrices with coefficients in $\Bbbk$ (see [Rad90]) to derive an example of a quasitriangular Hopf $\mathrm{GL}_n(\Bbbk)$-coalgebra.

Definition-Proposition 4.1. *For $\alpha = (\alpha_{i,j}) \in \mathrm{GL}_n(\Bbbk)$, let $\mathcal{A}_n^\alpha$ be the $\mathbb{C}$-algebra generated g, $x_1, \dots x_n$, $y_1, \dots, y_n$, subject to the following relations*

$$g^2 = 1, \quad x_1^2 = \cdots = x_n^2 = 0, \quad g x_i = -x_i g, \quad x_i x_j = -x_j x_i, \tag{4.1}$$

$$y_1^2 = \cdots = y_n^2 = 0, \quad g y_i = -y_i g, \quad y_i y_j = -y_j y_i, \tag{4.2}$$

$$x_i y_j - y_j x_i = (\delta_{i,j} - \alpha_{j,i})\, g, \tag{4.3}$$

where $1 \leq i, j \leq n$. The family $\mathcal{A}_n = \{\mathcal{A}_n^\alpha\}_{\alpha \in \mathrm{GL}_n(\Bbbk)}$ has a structure of a crossed Hopf $\mathrm{GL}_n(\Bbbk)$-coalgebra given, for any $\alpha = (\alpha_{i,j}) \in \mathrm{GL}_n(\Bbbk)$, $\beta = (\beta_{i,j}) \in \mathrm{GL}_n(\Bbbk)$, *and* $1 \leq i \leq n$, *by*

$$\Delta_{\alpha,\beta}(g) = g \otimes g, \quad \varepsilon(g) = 1, \quad S_\alpha(g) = g, \tag{4.4}$$

$$\Delta_{\alpha,\beta}(x_i) = 1 \otimes x_i + \sum_{k=1}^{n} \beta_{k,i}\, x_k \otimes g, \quad \varepsilon(x_i) = 0, \quad S_\alpha(x_i) = \sum_{k=1}^{n} \alpha_{k,i}\, gx_k, \tag{4.5}$$

$$\Delta_{\alpha,\beta}(y_i) = y_i \otimes 1 + g \otimes y_i, \quad \varepsilon(y_i) = 0, \quad S_\alpha(y_i) = -hy_i, \tag{4.6}$$

$$\varphi_\alpha(g) = g, \quad \varphi_\alpha(x_i) = \sum_{k=1}^{n} \alpha_{k,i}\, x_k, \quad \varphi_\alpha(y_i) = \sum_{k=1}^{n} \tilde{\alpha}_{k,i}\, y_k, \tag{4.7}$$

where $(\tilde{\alpha}_{i,j}) = \alpha^{-1}$. *Moreover* $\mathcal{A}_n$ *is quasitriangular with R-matrix given, for any* $\alpha, \beta \in \mathrm{GL}_n(\Bbbk)$, *by*

$$R_{\alpha,\beta} = \frac{1}{2} \sum_{S \subseteq [n]} x_S \otimes y_S + x_S \otimes gy_S + gx_S \otimes y_S + gx_S \otimes gy_S.$$

Here $[n] = \{1, \dots n\}$, $x_\emptyset = 1$, $y_\emptyset = 1$, *and, for a nonempty subset* S *of* $[n]$, *we let* $x_S = x_{i_1} \cdots x_{i_s}$ *and* $y_S = y_{i_1} \cdots y_{i_s}$ *where* $i_1 < \cdots < i_s$ *are the elements of* S.

Remark 4.2. From relations (4.3), it can be shown that the algebras $\mathcal{A}_n^\alpha$ and $\mathcal{A}_n^\beta$ are in general not isomorphic when $\alpha, \beta \in \mathrm{GL}_n(\Bbbk)$ are such that $\alpha \neq \beta$.

Proof. Let A_n be the $\Bbbk$-algebra generated by $g, x_1, \dots, x_n$ which satisfy the relations (4.1). The algebra A_n is 2^{n+1}-dimensional and has a Hopf algebra structure given by

$$\begin{aligned} &\Delta(g) = g \otimes g, && \varepsilon(g) = 1, && S(g) = g, \\ &\Delta(x_i) = x_i \otimes g + 1 \otimes x_i, && \varepsilon(x_i) = 0, && S(x_i) = gx_i. \end{aligned}$$

Radford [Rad90] showed that the group of Hopf automorphisms of A_n is isomorphic to the group $\mathrm{GL}_n(\Bbbk)$ of invertible $n \times n$-matrices with coefficients in $\Bbbk$. This group automorphism $\phi\colon \mathrm{GL}_n(\Bbbk) \to \mathrm{Aut}_{\mathrm{Hopf}}(A_n)$ is given, for any $\alpha = (\alpha_{i,j}) \in \mathrm{GL}_n(\Bbbk)$, by

$$\phi_\alpha(g) = g \quad \text{and} \quad \phi_\alpha(x_i) = \sum_{k=1}^{n} \alpha_{k,i}\, x_k.$$

The Hopf algebra $B_n = A_n^{\mathrm{cop}}$ is the $\Bbbk$-algebra generated by the symbols $h, y_1, \dots, y_n$ which satisfy the relations $h^2 = 1$ and (4.2) and its Hopf algebra structure is given by

$$\begin{aligned} &\Delta(h) = h \otimes h, && \varepsilon(h) = 1, && S(h) = h, \\ &\Delta(y_i) = y_i \otimes 1 + h \otimes y_i, && \varepsilon(y_i) = 0, && S(y_i) = -hy_i. \end{aligned}$$

Let us denote the cardinality of a set T by $|T|$. The elements $g^k x_S$ (resp. $h^k y_S$), where $k \in \{0, 1\}$ and $S \subseteq [n]$, form a basis for A_n (resp. B_n). Since Δ is multiplicative, it follows that

$$\Delta(g^k x_S) = \sum_{T \subseteq S} \lambda_{T,S}\; g^k x_T \otimes g^{k+|T|} x_{S \setminus T} \quad \text{and} \tag{4.8}$$

$$\Delta(h^k y_S) = \sum_{T \subseteq S} \lambda_{T,S}\; h^{k+|T|} y_{S \setminus T} \otimes h^k y_T, \tag{4.9}$$

where $\lambda_{T,S} = \pm 1$ and $\lambda_{\emptyset,S} = 1 = \lambda_{S,S}$.

By Section 2.1, there exists a (unique) Hopf pairing $\sigma\colon A_n \times B_n \to \Bbbk$ such that, for any $1 \leq i, j \leq n$,

$$\sigma(g,h) = -1, \quad \sigma(g, y_j) = \sigma(x_i, h) = 0 \quad \text{and} \quad \sigma(x_i, y_j) = \delta_{i,j}.$$

Using (4.8) and (4.9), one gets (by induction on $|S|$) that

$$\sigma(g^k x_S, h^l y_T) = (-1)^{kl}\, \delta_{S,T}$$

for any $k, l \in \{0,1\}$ and $S, T \subseteq [n]$, where $\delta_{S,S} = 1$ and $\delta_{S,T}$ if $S \neq T$. Set $z_0 = (1+h)/2$ and $z_1 = (1-h)/2$. The elements $z_k y_S$, where $k \in \{0,1\}$ and $S \subseteq [n]$, form a basis for B_n such that

$$\sigma(g^k x_S, z_l y_T) = \delta_{k,l}\, \delta_{S,T} \tag{4.10}$$

for any $k, l \in \{0,1\}$ and $S, T \subseteq [n]$. Therefore the pairing σ is non-degenerate. Note that this implies that $A_n^* \cong A_n$ as a Hopf algebra.

By Theorem 2.6, we get a quasitriangular Hopf $\mathrm{GL}_n(\Bbbk)$-coalgebra $D(A_n, B_n; \sigma, \phi)$. For any $\alpha = (\alpha_{i,j}) \in \mathrm{GL}_n(\Bbbk)$, $D(A_n, B_n; \sigma, \phi_\alpha)$ is the algebra generated by g, h, $x_1, \dots x_n$, $y_1, \dots, y_n$, subject to the relations $h^2 = 1$, (4.1), (4.2) and the following relations

$$gh = hg, \quad gy_j = -y_j g, \quad hx_i = x_i h, \tag{4.11}$$

$$x_i y_j - y_j x_i = \delta_{i,j} h - \alpha_{j,i} g. \tag{4.12}$$

Indeed $D(A_n, B_n; \sigma, \phi_\alpha)$ is the free algebra generated by the algebras A_n and B_n with cross relation (2.5). Further, it suffices to require the cross relations (2.5) for $(1 \otimes b) \cdot (a \otimes 1)$ with $a = g, x_i$ and $b = h, y_j$. To simplify the notations, we identify of a with $a \otimes 1$ and of b with $1 \otimes b$ (recall that these natural maps $A_n \hookrightarrow D(A_n, B_n; \sigma, \phi_\alpha)$ and $B_n \hookrightarrow D(A_n, B_n; \sigma, \phi_\alpha)$ are algebra monomorphisms). For example, let $a = x_i$ and $b = y_j$. Since $\sigma(x_i, 1) = \sigma(g, y_j) = \sigma(x_i, h) = \sigma(1, y_j) = 0$, relation (2.5) gives

$$y_j x_i = \sigma(\phi_\alpha(x_i), y_j h) \sigma(g, 1)\, g \cdot 1 + \sigma(1, h) \sigma(g, 1) x_i \cdot y_j + \sigma(1, h) \sigma(x_i, y_j) 1 \cdot h.$$

Inserting the values $\sigma(g, 1) = \sigma(1, h) = 1$, $\sigma(x_i, y_j) = \delta_{i,j}$, and $\sigma(\phi_\alpha(x_i), y_j h) = -\alpha_{j,i}$, we get (4.12).

From Theorem 2.3, we obtain that the comultiplication $\Delta_{\alpha,\beta}$, the counit ε, the antipode S_α, and the crossing φ_α of $D(A_n, B_n; \sigma, \phi_\alpha)$ are given by

$$\Delta_{\alpha,\beta}(g) = g \otimes g, \quad \Delta_{\alpha,\beta}(h) = h \otimes h, \tag{4.13}$$

$$\Delta_{\alpha,\beta}(x_i) = 1 \otimes x_i + \sum_{k=1}^{n} \beta_{k,i}\, x_k \otimes g, \quad \Delta_{\alpha,\beta}(y_i) = y_i \otimes 1 + h \otimes y_i, \tag{4.14}$$

$$\varepsilon(g) = \varepsilon(h) = 1, \quad \varepsilon(x_i) = \varepsilon(y_i) = 0, \quad S_\alpha(g) = g, \tag{4.15}$$

$$S_\alpha(h) = h, \quad S_\alpha(x_i) = \sum_{k=1}^{n} \alpha_{k,i}\, g x_k, \quad S_\alpha(y_i) = -h y_i, \tag{4.16}$$

$$\varphi_\alpha(g) = g, \quad \varphi_\alpha(h) = h, \quad \varphi_\alpha(x_i) = \sum_{k=1}^{n} \alpha_{k,i}\, x_k, \quad \varphi_\alpha(y_i) = \sum_{k=1}^{n} \tilde{\alpha}_{k,i}\, y_k, \tag{4.17}$$

where $(\tilde{\alpha}_{i,j}) = \alpha^{-1}$.

For any $\alpha \in \mathrm{GL}_n(\Bbbk)$, let I_α be the ideal of $D(A_n, B_n; \sigma, \phi_\alpha)$ generated by $g - h$. Using the above description of the structure maps of $D(A_n, B_n; \sigma, \phi)$, we get that $I = \{I_\alpha\}_{\alpha \in \pi}$ is a crossed Hopf $\mathrm{GL}_n(\Bbbk)$-coideal of $D(A_n, B_n; \sigma, \phi)$. The quotient $D(A_n, B_n; \sigma, \phi)/I = \{D(A_n, B_n; \sigma, \phi_\alpha)/I_\alpha\}_{\alpha \in \mathrm{GL}_n(\Bbbk)}$ is is precisely

$\mathcal{A}_n = \{\mathcal{A}_n^\alpha\}_{\alpha \in \mathrm{GL}_n(\Bbbk)}$ and so the latter has a quasitriangular Hopf $\mathrm{GL}_n(\Bbbk)$-coalgebra structure which can be described by replacing h with g in (4.13)-(4.17).

Finally, the R-matrix of $\mathcal{A}_n$ is obtained as the image under the projection maps $D(A_n, B_n; \sigma, \phi_\alpha) \overset{p_\alpha}{\to} D(A_n, B_n; \sigma, \phi_\alpha)/I_\alpha = \mathcal{A}_n^\alpha$ of the R-matrix of $D(A_n, B_n; \sigma, \phi)$, that is, using (4.10),

$$\begin{aligned} R_{\alpha,\beta} &= \sum_{S \subseteq [n]} p_\alpha(x_S) \otimes p_\beta(z_0 y_S) + p_\alpha(g x_S) \otimes p_\beta(z_1 y_S) \\ &= \sum_{S \subseteq [n]} x_S \otimes (\frac{1+g}{2}) y_S + g x_S \otimes (\frac{1-g}{2}) y_S \\ &= \frac{1}{2} \sum_{S \subseteq [n]} x_S \otimes y_S + x_S \otimes g y_S + g x_S \otimes y_S + g x_S \otimes g y_S. \end{aligned}$$

□

5. Graded quantum groups

Let $\mathfrak{g}$ be a finite-dimensional complex simple Lie algebra of rank l with Cartan matrix $(a_{i,j})$. We let d_i be the coprime integers such that the matrix $(d_i a_{i,j})$ is symmetric. Let q be a fixed non-zero complex number and let $q_i = q^{d_i}$. Suppose that $q_i^2 \neq 1$ for $i = 1, 2, \ldots, l$.

Definition-Proposition 5.1. *Set $\pi = (\mathbb{C}^*)^l$. For $\alpha = (\alpha_1, \ldots, \alpha_l) \in \pi$, let $U_q^\alpha(\mathfrak{g})$ be the $\mathbb{C}$-algebra generated by $K_i^{\pm 1}$, E_i, F_i, $1 \leq i \leq l$, subject to the following defining relations:*

$$K_i K_j = K_j K_i, \quad K_i K_i^{-1} = K_i^{-1} K_i = 1, \tag{5.1}$$

$$K_i E_j = q_i^{a_{i,j}} E_j K_i, \tag{5.2}$$

$$K_i F_j = q_i^{-a_{i,j}} F_j K_i, \tag{5.3}$$

$$E_i F_j - F_j E_i = \delta_{i,j} \frac{\alpha_i K_i - K_i^{-1}}{q_i - q_i^{-1}} \tag{5.4}$$

$$\sum_{r=0}^{1-a_{i,j}} (-1)^r \begin{bmatrix} 1-a_{i,j} \\ r \end{bmatrix}_{q_i} E_i^{1-a_{i,j}-r} E_j E_i^r = 0 \quad \text{if} \quad i \neq j, \tag{5.5}$$

$$\sum_{r=0}^{1-a_{i,j}} (-1)^r \begin{bmatrix} 1-a_{i,j} \\ r \end{bmatrix}_{q_i} F_i^{1-a_{i,j}-r} F_j F_i^r = 0 \quad \text{if} \quad i \neq j. \tag{5.6}$$

The family $U_q^\pi(\mathfrak{g}) = \{U_q^\alpha(\mathfrak{g})\}_{\alpha \in \pi}$ has a structure of a crossed Hopf π-coalgebra given, for $\alpha = (\alpha_1, \ldots, \alpha_l) \in \pi$, $\beta = (\beta_1, \ldots, \beta_l) \in \pi$ and $1 \leq i \leq l$, by

$$\begin{aligned} &\Delta_{\alpha,\beta}(K_i) = K_i \otimes K_i, \\ &\Delta_{\alpha,\beta}(E_i) = \beta_i E_i \otimes K_i + 1 \otimes E_i, \\ &\Delta_{\alpha,\beta}(F_i) = F_i \otimes 1 + K_i^{-1} \otimes F_i, \\ &\varepsilon(K_i) = 1, \quad \varepsilon(E_i) = \varepsilon(F_i) = 0, \\ &S_\alpha(K_i) = K_i^{-1}, \quad S_\alpha(E_i) = -\alpha_i E_i K_i^{-1}, \quad S_\alpha(F_i) = -K_i F_i, \\ &\varphi_\alpha(K_i) = K_i, \quad \varphi_\alpha(E_i) = \alpha_i E_i, \quad \varphi_\alpha(F_i) = \alpha_i^{-1} F_i. \end{aligned}$$

Remark 5.2. Note that $(U_q^1(\mathfrak{g}),\Delta_{1,1},\varepsilon,S_1)$ is the usual quantum group $U_q(\mathfrak{g})$.

Proof. Let U_+ be the $\mathbb{C}$-algebra generated by E_i, $K_i^{\pm 1}$, $1\leq i\leq l$, subject to the relations (5.1), (5.2) and (5.5). Let U_- be the $\mathbb{C}$-algebra generated by F_i, $K_i'^{\pm 1}$, $1\leq i\leq l$, subject to the relations (5.1), (5.3) and (5.6) where one has to replace K_i with K_i'. The algebras U_+ and U_- have a Hopf algebra structure given by

$$\begin{aligned}
&\Delta(K_i)=K_i\otimes K_i, \quad \Delta(E_i)=E_i\otimes K_i+1\otimes E_i,\\
&\varepsilon(K_i)=1, \quad \varepsilon(E_i)=0, \quad S(K_i)=K_i^{-1}, \quad S(E_i)=-E_iK_i^{-1},\\
&\Delta(K_i')=K_i'\otimes K_i', \quad \Delta(F_i)=F_i\otimes 1+K_i'^{-1}\otimes F_i,\\
&\varepsilon(K_i')=1, \quad \varepsilon(F_i)=0, \quad S(K_i)=K_i^{-1}, \quad S(F_i)=-K_i'F_i.
\end{aligned}$$

Using the method described in Section 2.1, it can be verified that there exists a (unique) Hopf pairing $\sigma\colon U_+\times U_-\to\mathbb{C}$ such that

$$\sigma(E_i,F_j)=\frac{\delta_{i,j}}{q_i-q_i^{-1}}, \quad \sigma(E_i,K_j')=\sigma(K_i,F_j)=0, \quad \sigma(K_i,K_j')=q_i^{a_{i,j}}=q_j^{a_{j,i}}.$$

Let $\phi:\pi\to\mathrm{Aut}_{\mathrm{Hopf}}(U_+)$ and $\psi:\pi\to\mathrm{Aut}_{\mathrm{Hopf}}(U_-)$ defined, for $\beta=(\beta_1,\dots,\beta_l)\in\pi$ and $1\leq i\leq l$, by

$$\phi_\beta(K_i)=K_i, \quad \phi_\beta(E_i)=\beta_i\,E_i, \quad \psi_\beta(K_i')=K_i', \quad \psi_\beta(F_i)=\beta_i^{-1}\,F_i.$$

It is straightforward to verify that ψ is (σ,ϕ)-compatible. By Lemma 2.4, we can consider the crossed Hopf π-coalgebra $D(U_+,U_-;\sigma,\phi)=\{D(U_+,U_-;\sigma,\phi_\alpha)\}_{\alpha\in\pi}$.

Now, for any $\alpha\in\pi$, $D(U_+,U_-;\sigma,\phi_\alpha)$ is the algebra generated by $K_i^{\pm1}$, $K_i'^{\pm1}$, E_i, F_i, where $1\leq i\leq l$, subject to the relations (5.1), (5.2), (5.5), the relations (5.1), (5.3), (5.6) where one has to replace K_i with K_i', and the following relations

$$K_iK_j'=K_j'K_i, \quad K_iF_j=q_i^{-a_{i,j}}F_jK_i, \quad K_i'E_j=q_i^{a_{i,j}}E_jK_i', \tag{5.7}$$

$$E_iF_j-F_jE_i=\delta_{i,j}\frac{\alpha_iK_i-K_i'^{-1}}{q_i-q_i^{-1}}. \tag{5.8}$$

Indeed $D(U_+,U_-;\sigma,\phi_\alpha)$ is the free algebra generated by the algebras U_+ and U_- with cross relation (2.5). Further, it suffices to require the cross relations (2.5) for $(1\otimes b)\cdot(a\otimes 1)$ with $a=K_i,E_i$ and $b=K_i',F_i$. To simplify the notations, we identify of a with $a\otimes 1$ and of b with $1\otimes b$ (recall that these natural maps $U_+\hookrightarrow D(U_+,U_-;\sigma,\phi_\alpha)$ and $U_-\hookrightarrow D(U_+,U_-;\sigma,\phi_\alpha)$ are algebra monomorphisms). For example, let $a=E_i$ and $b=F_j$. Since $\sigma(E_i,1)=\sigma(K_i,F_j)=\sigma(E_i,K_j'^{-1})=\sigma(1,F_j)=0$, relation (2.5) gives

$$F_jE_i=\sigma(\alpha_iE_i,S(F_j))\sigma(K_i,1)\,K_i+\sigma(1,K_j')\sigma(K_i,1)E_iF_j+\sigma(1,K_j')\sigma(E_i,F_j)\,K_j'^{-1}.$$

Inserting the values $\sigma(K_i,1)=\sigma(1,K_j')=1$, $\sigma(E_i,F_j)=\delta_{i,j}(q_i-q_i^{-1})^{-1}$ and $\sigma(E_i,S(F_j))=-\delta_{i,j}(q_i-q_i^{-1})^{-1}$, we get (5.8).

From Theorem 2.3, we obtain that the comultiplication $\Delta_{\alpha,\beta}$, the counit ε, the antipode S_α, and the crossing φ_α of $D(U_+,U_-;\sigma,\phi)$ are given, for $1\leq i\leq l$, by

$$\Delta_{\alpha,\beta}(K_i)=K_i\otimes K_i,\quad \Delta_{\alpha,\beta}(K'_i)=K'_i\otimes K'_i, \tag{5.9}$$

$$\Delta_{\alpha,\beta}(E_i)=\beta_i E_i\otimes K_i+1\otimes E_i,\quad \Delta_{\alpha,\beta}(F_i)=F_i\otimes 1+{K'_i}^{-1}\otimes F_i, \tag{5.10}$$

$$\varepsilon(K_i)=\varepsilon(K'_i)=1,\quad \varepsilon(E_i)=\varepsilon(F_i)=0,\quad S_\alpha(K_i)=K_i^{-1}, \tag{5.11}$$

$$S_\alpha(K'_i)={K'_i}^{-1},\quad S_\alpha(E_i)=-\alpha_i E_i K_i^{-1},\quad S_\alpha(F_i)=-K'_iF_i, \tag{5.12}$$

$$\varphi_\alpha(K_i)=K_i,\quad \varphi_\alpha(K'_i)=K'_i,\quad \varphi_\alpha(E_i)=\alpha_iE_i,\quad \varphi_\alpha(F_i)=\alpha_i^{-1}F_i. \tag{5.13}$$

Finally, for any $\alpha\in\pi$, let I_α be the ideal of $D(U_+,U_-;\sigma,\phi_\alpha)$ generated by $K_i-K'_i$ and $K_i^{-1}-{K'_i}^{-1}$, where $1\leq i\leq l$. Using the above description of the structure maps of $D(U_+,U_-;\sigma,\phi)$, we get that $I=\{I_\alpha\}_{\alpha\in\pi}$ is a crossed Hopf π-coideal of $D(U_+,U_-;\sigma,\phi)$. The quotient $D(U_+,U_-;\sigma,\phi)/I=\{D(U_+,U_-;\sigma,\phi_\alpha)/I_\alpha\}_{\alpha\in\pi}$ is precisely $U_q^\pi(\mathfrak{g})=\{U_q^\alpha\}_{\alpha\in\pi}$ and so the latter has a crossed Hopf π-coalgebra structure which can be described by replacing K'_i with K_i in (5.9)-(5.13). □

6. h-ADIC GRADED QUANTUM GROUPS

Let $\mathfrak{g}$ be a finite-dimensional complex simple Lie algebra of rank l with Cartan matrix $(a_{i,j})$. We let d_i be the coprime integers such that the matrix $(d_ia_{i,j})$ is symmetric.

Definition-Proposition 6.1. *Set $\pi=\mathbb{C}[[h]]^l$. For $\alpha=(\alpha_1,\dots,\alpha_l)\in\pi$, let $U_h^\alpha(\mathfrak{g})$ be the algebra over $\mathbb{C}[[h]]$ topologically generated by the elements H_i, E_i, F_i, $1\leq i\leq l$, subject to the following defining relations:*

$$[H_i,H_j]=0, \tag{6.1}$$

$$[H_i,E_j]=a_{ij}E_j, \tag{6.2}$$

$$[H_i,F_j]=-a_{ij}F_j, \tag{6.3}$$

$$[E_i,F_j]=\delta_{i,j}\frac{e^{d_ih\alpha_i}e^{d_ihH_i}-e^{-d_ihH_i}}{e^{d_ih}-e^{-d_ih}} \tag{6.4}$$

$$\sum_{r=0}^{1-a_{i,j}}(-1)^r\begin{bmatrix}1-a_{i,j}\\ r\end{bmatrix}_{e^{d_ih}}E_i^{1-a_{i,j}-r}E_jE_i^r=0\quad(i\neq j), \tag{6.5}$$

$$\sum_{r=0}^{1-a_{i,j}}(-1)^r\begin{bmatrix}1-a_{i,j}\\ r\end{bmatrix}_{e^{d_ih}}F_i^{1-a_{i,j}-r}F_jF_i^r=0\quad(i\neq j). \tag{6.6}$$

The family $U_q^\pi(\mathfrak{g})=\{U_q^\alpha(\mathfrak{g})\}_{\alpha\in\pi}$ has a structure of a crossed Hopf π-coalgebra given, for $\alpha=(\alpha_1,\dots,\alpha_l)\in\pi$, $\beta=(\beta_1,\dots,\beta_l)\in\pi$ and $1\leq i\leq l$, by

$$\Delta_{\alpha,\beta}(H_i)=H_i\otimes 1+1\otimes H_i,$$

$$\Delta_{\alpha,\beta}(E_i)=e^{d_ih\beta_i}E_i\otimes e^{d_ihH_i}+1\otimes E_i,$$

$$\Delta_{\alpha,\beta}(F_i)=F_i\otimes 1+e^{-d_ihH_i}\otimes F_i,$$

$$\varepsilon(H_i)=\varepsilon(E_i)=\varepsilon(F_i)=0,$$

$$S_\alpha(H_i)=-H_i,\quad S_\alpha(E_i)=-e^{d_ih\alpha_i}E_ie^{-d_ihH_i},\quad S_\alpha(F_i)=-e^{d_ihH_i}F_i,$$

$$\varphi_\alpha(H_i)=H_i,\quad \varphi_\alpha(E_i)=e^{d_ih\alpha_i}E_i,\quad \varphi_\alpha(F_i)=e^{-d_ih\alpha_i}F_i.$$

Remarks 6.2. (a) $(U_h^0(\mathfrak{g}), \Delta_{0,0}, \varepsilon, S_0)$ is the usual quantum group $U_h(\mathfrak{g})$.

(b) The element $e^{d_i h} - e^{-d_i h} \in \mathbb{C}[[h]]$ is not invertible in $\mathbb{C}[[h]]$, because the constant term is zero. But the expression oh the right hand side of (6.4) is a formal power series $\sigma_n p_n(H_i) h^n$ with certain polynomials $p_n(H_i)$, and so it is a well-defined element of the h-adic algebra generated by E_i, F_i, H_i.

Proof. Let U_+ be the h-adic algebra generated by H_i, E_i, $1 \leq i \leq l$, subject to the relations (6.1), (6.2) and (6.5). Let U_- be the h-adic algebra generated by H'_i, F_i, $1 \leq i \leq l$, subject to the relations (6.1), (6.3) and (6.6) where one has to replace H_i with H'_i. The algebras U_+ and U_- have a h-adic Hopf algebra structure given by

$$\begin{aligned}
&\Delta(H_i) = H_i \otimes 1 + 1 \otimes H_i, \quad \Delta(E_i) = E_i \otimes e^{d_i h H_i} + 1 \otimes E_i,\\
&\varepsilon(H_i) = \varepsilon(E_i) = 0, \quad S(H_i) = -H_i, \quad S(E_i) = -E_i e^{-d_i h H_i},\\
&\Delta(H'_i) = H'_i \otimes 1 + 1 \otimes H'_i, \quad \Delta(F_i) = F_i \otimes 1 + e^{-d_i h H'_i} \otimes F_i,\\
&\varepsilon(H'_i) = \varepsilon(F_i) = 0, \quad S(H'_i) = -H'_i, \quad S(F_i) = -e^{d_i h H'_i} F_i.
\end{aligned}$$

Let us consider the h-adic Hopf algebra $\widetilde{U}_- = \mathbb{C}[[h]]1 + hU_-$. The elements $\widetilde{H}'_i = hH'_i$ and $\widetilde{F}_i = hF_i$ belong to $\widetilde{U}_-$ and satisfy

$$[\widetilde{H}_i, \widetilde{F}_j] = -ha_{ij}\widetilde{F}_j, \quad \Delta(\widetilde{H}'_i) = \widetilde{H}'_i \otimes 1 + 1 \otimes \widetilde{H}'_i, \quad \Delta(\widetilde{F}_i) = \widetilde{F}_i \otimes 1 + e^{-d_i \widetilde{H}'_i} \otimes \widetilde{F}_i.$$

The element $e^{-d_i \widetilde{H}'_i} = 1 + \sum_{k \geq 1} \frac{1}{k!}(-d_i h)^k H_i^k$ is also in $\widetilde{U}_-$. Note that $e^{-d_i \widetilde{H}'_i}$ is not in the h-adic subalgebra of $\widetilde{U}_-$ generated by $\widetilde{H}'_i$.

Using the method described in Section 2.1, it can be verified that there exists a (unique) Hopf pairing $\sigma \colon U_+ \times \widetilde{U}_- \to \mathbb{C}[[h]]$ such that

$$\sigma(H_i, \widetilde{H}'_j) = d_i^{-1} a_{j,i}, \quad \sigma(H_i, \widetilde{F}_j) = \sigma(E_i, \widetilde{H}'_j) = 0, \quad \sigma(E_i, \widetilde{F}_j) = \frac{\delta_{i,j}\, h}{e^{d_i h} - e^{-d_i h}}.$$

Let $\phi : \pi \to \mathrm{Aut}_{\mathrm{Hopf}}(U_+)$ and $\psi : \pi \to \mathrm{Aut}_{\mathrm{Hopf}}(\widetilde{U}_-)$ defined, for $\alpha = (\alpha_1, \ldots, \alpha_l) \in \pi$ and $1 \leq i \leq l$, by

$$\phi_\alpha(H_i) = H_i, \quad \phi_\alpha(E_i) = e^{d_i h \alpha_i}\, E_i, \quad \psi_\alpha(\widetilde{H}'_i) = \widetilde{H}'_i, \quad \psi_\beta(\widetilde{F}_i) = e^{-d_i h \alpha_i}\, \widetilde{F}_i.$$

It is straightforward to verify that ψ is (σ, ϕ)-compatible. By the h-adic version of Lemma 2.4, we can consider the crossed h-adic Hopf π-coalgebra $D(U_+, \widetilde{U}_-; \sigma, \phi) = \{D(U_+, \widetilde{U}_-; \sigma, \phi_\alpha)\}_{\alpha \in \pi}$ whose structure can be explicitly described as in the proof of Proposition 5.1.

For any $\alpha \in \pi$, let I_α be the h-adic ideal of $D(U_+, \widetilde{U}_-; \sigma, \phi_\alpha)$ generated by $\widetilde{H}'_i - hH_i$ where $1 \leq i \leq l$. Using the description of the structure maps of $D(U_+, \widetilde{U}_-; \sigma, \phi_\alpha)$, we get that $I = \{I_\alpha\}_{\alpha \in \pi}$ is a crossed h-adic Hopf π-coideal of $D(U_+, U_-; \sigma, \phi)$. The quotient $D(U_+, \widetilde{U}_-; \sigma, \phi)/I = \{D(U_+, \widetilde{U}_-; \sigma, \phi_\alpha)/I_\alpha\}_{\alpha \in \pi}$ is precisely $U_h^\pi(\mathfrak{g}) = \{U_h^\alpha\}_{\alpha \in \pi}$ and so the latter has a crossed h-adic Hopf π-coalgebra structure. □

It is well-know (see, e.g., [KS97]) that the Hopf pairing $\sigma \colon U_+ \times \widetilde{U}_- \to \mathbb{C}[[h]]$ is non-degenerate and that, if $(e_i)_i$ and $(f_i)_i$ are dual basis of the vector spaces U_+ and $\widetilde{U}_-$ with respect to the form σ, then $\sum_i (e_i \otimes 1) \otimes (1 \otimes f_i)$ belongs to the h-adic completion $D(U_+, \widetilde{U}_-; \sigma, \phi_\alpha) \hat{\otimes} D(U_+, \widetilde{U}_-; \sigma, \phi_\beta)$. Therefore, by Theorem 2.10, the crossed h-adic Hopf π-coalgebra $D(U_+, \widetilde{U}_-; \sigma, \phi)$ is quasitriangular. Hence, as a quotient of $D(U_+, \widetilde{U}_-; \sigma, \phi)$, $U_h^\pi(\mathfrak{g})$ is also quasitriangular.

For example, when $\mathfrak{g} = \mathfrak{sl}_2$ and so $\pi = \mathbb{C}[[h]]$, we have that the R-matrix of $U_h^{\mathbb{C}[[h]]}(\mathfrak{sl}_2)$ is given, for any $\alpha, \beta \in \mathbb{C}[[h]]$, by

$$R_{\alpha,\beta} = e^{h(H \otimes H)/2} \sum_{n=0}^{\infty} R_n(h)\, E^n \otimes F^n,$$

where $R_n(h) = q^{n(n+1)/2} \frac{(1-q^{-2})^n}{[n]_q!}$ and $q = e^h$.

For $n \geq 1$, there exits a representation $\rho_n^\alpha : U_h^\alpha(\mathfrak{sl}_2) \to \mathrm{GL}(V_n^\alpha)$, where $V_n^\alpha = \mathbb{C}^n$ as a vector space, given on the standard basis $(e_i)_{1 \leq i \leq n}$ of $\mathbb{C}^n$ by

$$\rho_n^\alpha(H) e_i = (n - 2i + 1 - \frac{\alpha}{2})\, e_i,$$

$$\rho_n^\alpha(E) e_i = \begin{cases} e^{\frac{h\alpha}{2}}\, [n-i+1]_q\, e_{i-1} & \text{if } i > 1 \\ 0 & \text{if } i = 0 \end{cases},$$

$$\rho_n^\alpha(F) e_i = \begin{cases} [i]_q\, e_{i+1} & \text{if } i < n \\ 0 & \text{if } i = n \end{cases}.$$

Together with the quasitriangularity of $U_h^{\mathbb{C}[[h]]}(\mathfrak{sl}_2)$, this data leads in particular to a solution of the $\mathbb{C}[[h]]$-colored Yang-Baxter equation.

References

[Dae93] A. Van Daele, *Dual pairs of Hopf $*$-algebras*, Bull. London Math. Soc. **25** (1993), no. 3, 209–230.

[Dri87] V. G. Drinfeld, *Quantum groups*, Proc. I.C.M. Berkeley 1986, Vol. 1, 2 (Berkeley, Calif., 1986) (Providence, RI), Amer. Math. Soc., 1987, pp. 798–820.

[KRT97] C. Kassel, M. Rosso, and V. Turaev, *Quantum groups and knot invariants*, Panoramas et Synthèses [Panoramas and Syntheses], vol. 5, Société Mathématique de France, 1997.

[KS97] A. Klimyk and K. Schmudgen, *Quantum Groups and Their Representations*, Springer-Verlag, Berlin, New York, 1997.

[Oht93] T. Ohtsuki, *Colored ribbon Hopf algebras and universal invariants of framed links*, J. Knot Theory and Its Rami. **2** (1993), no. 2, 211–232.

[Rad90] D. E. Radford, *The group of automorphisms of a semisimple Hopf algebra over a field of characteristic* 0 *is finite*, Amer. J. Math. **112** (1990), no. 2, 331–357.

[Tur00] V. Turaev, *Homotopy field theory in dimension* 3 *and group-categories*, preprint GT/0005291, 2000.

[Vir01] A. Virelizier, *Algèbres de Hopf graduées et fibrés plats sur les* 3*-variétés*, Ph.D. thesis, 2001.

[Vir02a] ______, *Hopf group-coalgebras*, J. Pure Appl. Algebra **171** (2002), no. 1, 75–122.

[Vir02b] ______, *Involutory Hopf algebras and flat bundles over 3-manifolds*, Prepublication, 2002.

[Zun02] M. Zunino, *Double construction for crossed Hopf coalgebras*, math.QA/0212192, 2002.

Département des Sciences Mathématiques, Université Montpellier II, Case Courrier 051, Place Eugène Bataillon, 34095 Montpellier Cedex 5, France

E-mail address: `virelizi@math.univ-montp2.fr`